



## ON EQUIVALENCE OF TWO INTEGRABILITY METHODS

H. N. ÖZGEN

*Received 17 November, 2015*

**Abstract.** In this paper, we introduce the concept of  $|R, p|_k, k \geq 1$  integrability of improper integrals and by this definition we prove a theorem, that generalizes a theorem of Orhan [3].

**2010 Mathematics Subject Classification:** 26D15; 40F05; 40D25; 40A10; 35A23

**Keywords:** Cesàro and Riesz method, improper integrals, divergent integrals, inequalities for integrals

### 1. INTRODUCTION

Throughout this paper we assume that  $f$  is a real valued function which is continuous on  $[0, \infty)$  and  $s(x) = \int_0^x f(t)dt$ . By  $\sigma(x)$ , we denote the Cesàro mean of  $s(x)$ . The integral  $\int_0^\infty f(t)dt$  is said to be integrable  $|C, 1|_k, k \geq 1$ , in the sense of Flett [2], if

$$\int_0^\infty x^{k-1} |\sigma'(x)|^k dx = \int_0^\infty \frac{|v(x)|^k}{x} dx \quad (1.1)$$

is convergent. Here,  $v(x) = \frac{1}{x} \int_0^x t f(t)dt$  is called a generator of the integral  $\int_0^\infty f(t)dt$ . Let  $p$  be a real valued, non-decreasing function on  $[0, \infty)$  such that

$$P(x) = \int_0^x p(t)dt, p(x) \neq 0, p(0) = 0.$$

The Riesz mean of  $s(x)$  is defined by

$$\sigma_p(x) = \frac{1}{P(x)} \int_0^x p(t)s(t)dt.$$

We say that the integral  $\int_0^\infty f(t)dt$  is integrable  $|R, p|_k, k \geq 1$ , if

$$\int_0^\infty x^{k-1} |\sigma_p'(x)|^k dx \quad (1.2)$$

is convergent. In the special case if we take  $p(x) = 1$  for all values of  $x$ , then  $|R, p|_k$  integrability reduces to  $|C, 1|_k$  integrability of improper integrals.

Given any functions  $f, g$ , it is customary to write  $g(x) = O(f(x))$ , if there exist  $\eta$

and  $N$ , for every  $x > N$ ,  $|\frac{g(x)}{f(x)}| \leq \eta$ .

The difference between  $s(x)$  and its  $n$ th weighted mean  $\sigma_p(x)$ , which is called the weighted Kronecker identity, is given by the identity

$$s(x) - \sigma_p(x) = v_p(x), \quad (1.3)$$

where

$$v_p(x) = \frac{1}{P(x)} \int_0^x P(u) f(u) du.$$

We note that if we take  $p(x) = 1$ , for all values of  $x$  then we have the following identity (see [1])

$$s(x) - \sigma(x) = v(x).$$

Since

$$\sigma'_p(x) = \frac{p(x)}{P(x)} v_p(x),$$

condition (1.3) can be rewritten as

$$s(x) = v_p(x) + \int_0^x \frac{p(u)}{P(u)} v_p(u) du. \quad (1.4)$$

In view of the identity (1.4), the function  $v_p(x)$  is called the generator function of  $s(x)$ .

Condition (1.1) can also be written as

$$\int_0^\infty x^{k-1} \left( \frac{P(x)}{P(x)} \right)^k |v_p(x)|^k dx \quad (1.5)$$

is convergent. We note that for infinite series, an analogous definition was introduced by Orhan [3]. Using this definition, Orhan [3] proved the following theorem dealing with  $|R, p_n|_k$  and  $|R, q_n|_k$  summability methods.

**Theorem 1.** *The  $|R, p_n|_k, (k \geq 1)$  summability implies the  $|R, q_n|_k, (k \geq 1)$  summability provided that*

$$n q_n = O(Q_n), \quad (1.6)$$

$$P_n = O(np_n), \quad (1.7)$$

$$Q_n = O(nq_n). \quad (1.8)$$

## 2. MAIN RESULT

The aim of this paper is to state Orhan's theorem for  $|R, p|_k$  and  $|R, q|_k$  integrability of improper integrals. Now we shall prove the following theorem.

**Theorem 2.** *Let  $p$  and  $q$  be real valued, non-decreasing functions on  $[0, \infty)$  such that as  $x \rightarrow \infty$*

$$xq(x) = O(Q(x)), \quad (2.1)$$

$$P(x) = O(xp(x)), \quad (2.2)$$

$$Q(x) = O(xq(x)). \quad (2.3)$$

*If  $\int_0^\infty f(t)dt$  is integrable  $|R, p|_k$ , then it is also integrable  $|R, q|_k, k \geq 1$ .*

## 3. PROOF OF THE THEOREM

Let  $\sigma_p(x)$  and  $\sigma_q(x)$  be the functions of  $(R, p)$  and  $(R, q)$  means of the integral  $\int_0^\infty f(t)dt$ . Since  $\int_0^\infty f(t)dt$  is integrable  $|R, p|_k$ , we can write

$$\int_0^\infty x^{k-1} \left( \frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx$$

is convergent. Differentiating the equation (1.4), we have

$$f(x) = v'_p(x) + \frac{p(x)}{P(x)} v_p(x).$$

By definition, we obtain

$$\sigma_q(x) = \frac{1}{Q(x)} \int_0^x q(t)s(t)dt = \frac{1}{Q(x)} \int_0^x (Q(x) - Q(t))f(t)dt$$

and

$$\begin{aligned} \sigma'_q(x) &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t)f(t)dt \\ &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \left[ v'_p(t) + \frac{p(t)}{P(t)} v_p(t) \right] dt \\ &= \frac{q(x)}{Q^2(x)} \int_0^x Q(t)v'_p(t)dt + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t)dt. \end{aligned}$$

Integrating by parts of the first statement, we have

$$\begin{aligned} \sigma'_q(x) &= \frac{q(x)}{Q^2(x)} \left[ Q(x)v_p(x) - \int_0^x q(t)v_p(t)dt \right] + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t)dt \\ &= \frac{q(x)}{Q(x)} v_p(x) + \frac{q(x)}{Q^2(x)} \int_0^x Q(t) \frac{p(t)}{P(t)} v_p(t)dt - \frac{q(x)}{Q^2(x)} \int_0^x q(t)v_p(t)dt \\ &= \sigma_{q,1}(x) + \sigma_{q,2}(x) + \sigma_{q,3}(x), \text{ say.} \end{aligned}$$

To complete the proof of the theorem, it is sufficient to show that

$$\int_0^m x^{k-1} |\sigma_{q,r}(x)|^k dx = O(1) \text{ as } m \rightarrow \infty, \text{ for } r = 1, 2, 3.$$

Using conditions (2.1) and (2.2), we have

$$\begin{aligned} \int_0^m x^{k-1} |\sigma_{q,1}(x)|^k dx &= \int_0^m x^{k-1} \left| \frac{q(x)}{Q(x)} v_p(x) \right|^k dx \\ &= \int_0^m x^{k-1} \left( \frac{q(x)}{Q(x)} \right)^k |v_p(x)|^k dx \\ &= O(1) \int_0^m x^{k-1} \left( \frac{p(x)}{P(x)} \right)^k |v_p(x)|^k dx \\ &= O(1) \int_0^m x^{k-1} |\sigma'_p(x)|^k dx \\ &= O(1) \text{ as } m \rightarrow \infty \end{aligned}$$

by virtue of the hypotheses of Theorem 2.

Applying Hölder's inequality with  $k > 1$ , we get

$$\begin{aligned} \int_0^m x^{k-1} |\sigma_{q,2}(x)|^k dx &= \\ &= O(1) \int_0^m x^{k-1} \left( \frac{q(x)}{Q^2(x)} \right)^k \left( \int_0^x \frac{Q(t)p(t)}{P(t)} |v_p(t)| dt \right)^k dx \\ &= O(1) \int_0^m \frac{q(x)}{Q^{k+1}(x)} \left( \int_0^x \frac{Q(t)p(t)}{P(t)} |v_p(t)| dt \right)^k dx \\ &= O(1) \int_0^m \frac{q(x)}{Q^2(x)} \left( \int_0^x \left( \frac{Q(t)}{q(t)} \right)^k q(t) \left( \frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \right) \\ &\quad \times \left( \frac{1}{Q(x)} \int_0^x q(t) dt \right)^{k-1} dx \\ &= O(1) \int_0^m t^k q(t) \left( \frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \int_t^m \frac{q(x)}{Q^2(x)} dx \\ &= O(1) \int_0^m t^k \frac{q(t)}{Q(t)} \left( \frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \\ &= O(1) \int_0^m t^{k-1} \left( \frac{p(t)}{P(t)} \right)^k |v_p(t)|^k dt \\ &= O(1) \int_0^m t^{k-1} |\sigma'_p(t)|^k dt \end{aligned}$$

$$= O(1) \text{ as } m \rightarrow \infty$$

by virtue of the hypotheses of Theorem 2.

Finally, again by Hölder's inequality with  $k > 1$ , we have

$$\begin{aligned} \int_0^m x^{k-1} |\sigma_{q,3}(x)|^k dx &= O(1) \int_0^m x^{k-1} \left( \frac{q(x)}{Q^2(x)} \right)^k \left( \int_0^x q(t) |v_p(t)|^k dt \right)^k dx \\ &= O(1) \int_0^m \frac{q(x)}{Q^2(x)} \left( \int_0^x q(t) |v_p(t)|^k dt \right) \\ &\quad x \left( \frac{1}{Q(x)} \int_0^x q(t) dt \right)^{k-1} dx \\ &= O(1) \int_0^m q(t) |v_p(t)|^k dt \int_t^m \frac{q(x)}{Q^2(x)} dx \\ &= O(1) \int_0^m \frac{q(t)}{Q(t)} |v_p(t)|^k dt \\ &= O(1) \text{ as } m \rightarrow \infty \end{aligned}$$

by virtue of the hypotheses of Theorem 2.

This completes the proof of the theorem.

**Theorem 3.** Let  $p$  and  $q$  be real valued, non-decreasing functions on  $[0, \infty)$  such that as  $x \rightarrow \infty$

$$xp(x) = O(P(x)), \quad (3.1)$$

$$Q(x) = O(xq(x)), \quad (3.2)$$

$$P(x) = O(xp(x)). \quad (3.3)$$

If  $\int_0^\infty f(t)dt$  is integrable  $|R, q|_k$ , then it is also integrable  $|R, p|_k, k \geq 1$ .

*Proof.* In Theorem 2 if we take  $p(x) = q(x)$  and  $q(x) = p(x)$ , then we get Theorem 3.  $\square$

**Theorem 4.** Let  $p$  and  $q$  be real valued, non-decreasing functions on  $[0, \infty)$  such that as  $x \rightarrow \infty$

$$xp(x) = O(P(x)), \quad (3.4)$$

$$P(x) = O(xp(x)), \quad (3.5)$$

$$xq(x) = O(Q(x)), \quad (3.6)$$

$$Q(x) = O(xq(x)). \quad (3.7)$$

Then the  $|R, p|_k$  integrability of  $\int_0^\infty f(t)dt$  is equivalent to the  $|R, q|_k$  integrability of  $\int_0^\infty f(t)dt$ , where  $k \geq 1$ .

## REFERENCES

- [1] I. Çanak and U. Totur, “A tauberian theorem for cesàro summability factors of integrals,” *Appl. Math. Lett.*, vol. 24, no. 3, pp. 234–245, 2011.
- [2] T. M. Flett, “On an extension of absolute summability and some theorems of littlewood and paley,” *Proc. London Math. Soc.*, vol. 7, pp. 113–141, 1957.
- [3] C. Orhan, “On equivalence of summability methods,” *Math. Slovaca.*, vol. 40, no. 2, pp. 171–175, 1990.

*Author's address*

**H. N. Özgen**

Department of Mathematics, Faculty of Education, Mersin University, TR-33169 Mersin, Turkey

*E-mail address:* nogduk@gmail.com