



AN EXISTENCE ETUDY FOR A TRIPLED SYSTEM WITH p –LAPLACIAN INVOLVING φ –CAPUTO DERIVATIVES

HAMID BEDDANI, MOUSTAFA BEDDANI, AND ZOUBIR DAHMANI

Received 05 January, 2022

Abstract. In this paper, we study the existence and uniqueness of solutions for a tripled system of fractional differential equations with nonlocal integro multi point boundary conditions by using the p –Laplacian operator and the φ –Caputo derivatives. The presented results are obtained by the two fixed point theorems of Banach and Krasnoselskii. An illustrative example is presented at the end to show the applicability of the obtained results. To the best of our knowledge, this is the first time where such problem is considered.

2010 *Mathematics Subject Classification:* 30C45; 39B72; 39B85

Keywords: p –Laplacian operator, φ –Caputo derivative, existence of solution, fixed point

1. INTRODUCTION

The fractional calculus has many significant roles in various scientific fields of research, see for instance [10, 20]. As applied results, the fractional order differential equations have attracted attention of several scientists in different fields of research [8, 19]. However, most of the published works have been achieved by using the fractional derivatives of type Riemann-Liouville, Hadamard, Katugampola, Atangana-Baleanu, Grunwald Letnikov and Caputo. The fractional derivatives of functions with respect to some other functions [14] are different from the others since their kernels appear in terms of other functions (called φ). Recently, some fractional differential results have been considered in [4–6, 11].

In most of the present articles, Schauder, Krasnoselskii, Darbo, or Monch theories have been used to prove existence of solutions of nonlinear fractional differential equations with some restrictive conditions [3, 18]. Some authors have worked on the solutions for fractional problems with p –Laplacian operators. We cite, for example [1, 2, 12, 16] where it has been studied nonlinear fractional equation with p –Laplacian operator for the solutions.

Here, we will mention some other research works for the reader. We begin by A. Devi, A. Kumar, D. Baleanu and A. Khan [2] where they worked on the stability results, for the following nonlinear FDEs involving Caputo derivatives of distinct

orders and Ψ_p Laplacian operator:

$$\begin{cases} {}^c \mathcal{D}^{r_1} \Psi_p [{}^c \mathcal{D}^{r_2} (u(t) - \sum_{i=1}^m v_i(t))] = -w(t, u(t)), t \in (0, 1) \\ \Psi_p [{}^c \mathcal{D}^{r_2} (u(t) - \sum_{i=1}^m v_i(t))] |_{t=0} = 0, \\ u(0) = \sum_{i=1}^m v_i(0), \\ u'(1) = \sum_{i=1}^m v_i'(1), \\ u^j(0) = \sum_{i=1}^m v_i^j(0), \text{ for } j = 2, 3, \dots, n-1, \end{cases}$$

where $0 < r_1 \leq 1, n-1 < r_2 \leq n, n \geq 4$, and v_i, w are continuous functions. ${}^c \mathcal{D}^{r_1}$ and ${}^c \mathcal{D}^{r_2}$ denotes the derivative of fractional order r_1 and r_2 in Caputo's sense, respectively, and $\Psi_p(z) = |z|^{p-2}z$ denotes the p -Laplacian operator and satisfies $\frac{1}{p} + \frac{1}{q} = 1, (\Psi_p)^{-1} = \Psi_q$.

We can also cite the paper of A. Mahdjouba et al. [17] where they have investigated the existence and multiplicity of positive solutions of the following problem:

$$\begin{cases} (\Psi_p [\mathcal{D}_{0+}^r (u(t))])' + a_1(t)f(u(\theta_1(t)), v(\theta_2(t))) = 0, 0 < t < 1, \\ (\Psi_{\bar{p}} [\mathcal{D}_{0+}^r (v(t))])' + a_2(t)f(u(\theta_1(t)), v(\theta_2(t))) = 0, 0 < t < 1, \\ \mathcal{D}_{0+}^r u(0) = u(0) = u'(0), \quad \mathcal{D}_{0+}^m u(1) = \gamma \mathcal{D}_{0+}^m u(\eta), \\ \mathcal{D}_{0+}^r v(0) = v(0) = v'(0), \quad \mathcal{D}_{0+}^m v(1) = \gamma \mathcal{D}_{0+}^m v(\eta), \end{cases}$$

where $\eta \in (0, 1), \gamma \in (0, \frac{1}{\eta^{r-m-1}})$, $\mathcal{D}_{0+}^r, \mathcal{D}_{0+}^m$, are the standard Riemann–Liouville fractional derivatives with $r \in (2, 3), m \in (1, 2)$ such that $r \geq m+1$, p -Laplacian operator is defined as $\Psi_p(z) = z|z|^{p-2}, p > 1$, and the functions $f, g \in C(\mathbb{R}^2, \mathbb{R})$.

Then, S. Etemad with his coauthors [9] have been concerned with the existence study for the following tripled impulsive fractional problem

$$\begin{cases} {}^c \mathcal{D}_{0+}^{\kappa_m} x_m(t) = f_m(t, x(t)), m = 1, 2, 3, \text{ and } t \in J' \\ x_m(a) = \Phi_m x, x'_m(a) = \Theta_m x, \\ \Delta x_m |_{t=t_k} = I_{m,k}(x(t_k)), \Delta x'_m |_{t=t_k} = \bar{I}_{m,k}(x(t_k)), \end{cases}$$

where $J = [a, b], J' = J - \{t_1, t_2, \dots, t_p\}, a = t_0 < t_1 < \dots < t_p < t_{p+1} = b$, ${}^c \mathcal{D}_{0+}^{\kappa_m}$, $m = 1, 2, 3$, are the Caputo fractional derivatives such that $\kappa_m \in (1, 2], f_m : J \times \mathbb{R}^3 \rightarrow \mathbb{R}, x(t) = (x_1(t), x_2(t), x_3(t)), I_{m,k}, \bar{I}_{m,k} : \mathbb{R}^3 \rightarrow \mathbb{R}, k = 1, 2, \dots, p$, are given functions, Φ_m, Θ_m are given operators, $\Delta x_m |_{t=t_k} = x(t_k^+) - x(t_k^-), \Delta x'_m |_{t=t_k} = x'(t_k^+) - x'(t_k^-)$, and

$$x(t_k^+) = \lim_{h \rightarrow 0^+} x_m(t_k + h), x(t_k^-) = \lim_{h \rightarrow 0^-} x_m(t_k + h).$$

In the present research work, we study the existence and uniqueness of solutions for the following problem:

$$\begin{cases} \mathcal{D}_{0+}^{r_{1m}; \Phi} \Psi_p [\mathcal{D}_{0+}^{r_{2m}; \Phi} (u_m(t) - I_{0+}^{\sigma; \Phi} G_m(t, u_1(t), u_2(t), u_3(t)))] \\ = H_m(t, u_1(t), u_2(t), u_3(t)), m = 1, 2, 3, \text{ and } t \in J = (0, 1] \\ \Psi_p [\mathcal{D}_{0+}^{r_{2m}; \Phi} (u_m(t) - I_{0+}^{\sigma; \Phi} G_m(t, u_1(t), u_2(t), u_3(t)))] |_{t=0} = 0, \\ u_m(0) = 0, \quad u_m(1) = \sum_{i=1}^n \lambda_{im} u_m(\zeta_{im}), \quad \zeta_{im} \in (0, 1] \\ \varphi(1) - \varphi(0) = K > 0. \end{cases} \quad (1.1)$$

Here, we take $\mathcal{D}_{0^+}^{r_{im};\varphi}, i = \overline{1, 2}, m = \overline{1, 2, 3}$ as the φ -Caputo fractional derivatives of orders $r_{im}, 0 < r_{1m} < 1 < r_{2m} < 2$, and $I_{0^+}^{\sigma;\varphi}, 0 < \sigma < r_{2m}$, the fractional integral of order $\sigma, \lambda_{im} \in \mathbb{R}_+^*$, and $\varphi : J \rightarrow \mathbb{R}$ is an increasing function such that $\varphi'(t) \neq 0$, and $\Psi_p(z) = |z|^{p-2}z$ denotes the p -Laplacian operator and satisfies $\frac{1}{p} + \frac{1}{q} = 1, (\Psi_p)^{-1} = \Psi_q$. For all $t \in J, G_m, H_m : J \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a given functions satisfying some assumptions that will be specified later.

2. φ -CAPUTO DERIVATIVES

In this section, we introduce some notations and definitions of φ -Caputo approach, for details, see [6, 14, 19].

Let $\varphi : J \rightarrow \mathbb{R}$ be an increasing function with $\varphi'(t) \neq 0$, for all $t \in J$.

And throughout the paper, let $C = C(J, \mathbb{R})$ denotes the Banach space of all continuous mappings from $[0, 1]$ to \mathbb{R} endowed with the norm $\|u\| = \sup_{t \in [0,1]} u(t)$. It is clear that the space $C \times C \times C$ endowed with the norm $\|(u_1, u_2, u_3)\| = \|u_1\| + \|u_2\| + \|u_3\|$ is a Banach space. We pose for all $r > 0$, and $t \in [0, 1], (t > s)$

$$\varphi_r(t, s) = \frac{\varphi'(s) (\varphi(t) - \varphi(s))^{r-1}}{\Gamma(r)}.$$

Definition 1. For $\alpha > 0$, the left-sided φ -Riemann Liouville fractional integral of order α for an integrable function $u : J \rightarrow \mathbb{R}$ with respect to another function $\varphi : J \rightarrow \mathbb{R}$ that is an increasing differentiable function such that $\varphi'(t) \neq 0$, for all $t \in J$ is defined as follows

$$I_{a^+}^{\alpha;\varphi} u(t) = \int_a^t \varphi_{\alpha}(t, s) u(s) ds, \tag{2.1}$$

where Γ is the gamma function. Note that equation (2.1) is reduced to the Riemann Liouville and Hadamard fractional integrals when $\varphi(t) = t$ and $\varphi(t) = \ln t$, respectively.

Definition 2. Let $n \in \mathbb{N}$ and let $\varphi, u \in C^n(J)$ be two functions such that φ is increasing and $\varphi'(t) \neq 0$, for all $t \in J$. The left-sided φ -Riemann Liouville fractional derivative of a function u of order α is defined by

$$\begin{aligned} \mathcal{D}_{a^+}^{\alpha;\varphi} u(t) &= \left(\frac{1}{\varphi'(t)} \frac{d}{dt} \right)^n I_{a^+}^{n-\alpha;\varphi} u(t) \\ &= \left(\frac{1}{\varphi'(t)} \frac{d}{dt} \right)^n \int_a^t \varphi_{n-\alpha}(t, s) u(s) ds, \end{aligned}$$

where $n = [\alpha] + 1$.

Definition 3. Let $n \in \mathbb{N}$ and let $\varphi, u \in C^n(J)$ be two functions such that φ is increasing and $\varphi'(t) \neq 0$, for all $t \in J$. The left-sided φ -Caputo fractional derivative of a function u of order α is defined by

$${}^c \mathcal{D}_{a^+}^{\alpha; \varphi} u(t) = I_{a^+}^{n-\alpha; \varphi} \left(\frac{1}{\varphi'(t)} \frac{d}{dt} \right)^n u(t),$$

where $n = [\alpha] + 1$ for $\alpha \notin \mathbb{N}$, $n = \alpha$ for $\alpha \in \mathbb{N}$.

To simplify notation, we will use the symbol

$$u_{\varphi}^{[n]}(t) = \left(\frac{1}{\varphi'(t)} \frac{d}{dt} \right)^n u(t).$$

From the definition, it is clear that

$${}^c \mathcal{D}_{a^+}^{\alpha; \varphi} u(t) = \begin{cases} \int_a^t \varphi_{n-\alpha}(t, s) u_{\varphi}^{[n]}(s) ds, & \text{if } \alpha \notin \mathbb{N}, \\ u_{\varphi}^{[n]}(t) & \text{if } \alpha \in \mathbb{N}. \end{cases} \quad (2.2)$$

This generalization (2.2) yields to the Caputo fractional derivative operator when $\varphi(t) = t$. Moreover, for $\varphi(t) = \ln t$, it gives the Caputo Hadamard fractional derivative.

2.1. Auxiliary Lemma

Lemma 1. Let $\alpha, \beta > 0$, and $u \in L^1(J)$. Then

$$I_{a^+}^{\alpha; \varphi} I_{a^+}^{\beta; \varphi} u(t) = I_{a^+}^{\alpha+\beta; \varphi} u(t), \quad \text{a.e. } t \in J.$$

In particular, if $u \in C(J)$, then $I_{a^+}^{\alpha; \varphi} I_{a^+}^{\beta; \varphi} u(t) = I_{a^+}^{\alpha+\beta; \varphi} u(t)$, $t \in J$.

Next, we recall the property describing the composition rules for fractional φ -integrals and φ -derivatives.

Lemma 2. Let $\alpha > 0$. The following holds:

If $u \in C([a, b])$, then

$${}^c \mathcal{D}_{a^+}^{\alpha; \varphi} I_{a^+}^{\alpha; \varphi} u(t) = u(t), \quad t \in [a, b].$$

If $u \in C^n(J)$, $n - 1 < \alpha < n$, then

$$I_{a^+}^{\alpha; \varphi c} \mathcal{D}_{a^+}^{\alpha; \varphi} u(t) = u(t) - \sum_{k=0}^{n-1} \frac{u_{\varphi}^{[k]}(a)}{k!} [\varphi(t) - \varphi(a)]^k,$$

for all $t \in [a, b]$. In particular, if $0 < \alpha < 1$, we have

$$I_{a^+}^{\alpha; \varphi c} \mathcal{D}_{a^+}^{\alpha; \varphi} u(t) = u(t) - u(a).$$

Lemma 3. Let $t > a$, $\alpha \geq 0$ and $\beta > 0$. Then

$$\bullet \quad I_{a^+}^{\alpha; \varphi} [\varphi(t) - \varphi(a)]^{\beta-1} = \frac{\Gamma(\beta)}{\Gamma(\beta+\alpha)} [\varphi(t) - \varphi(a)]^{\beta+\alpha-1},$$

- ${}^c \mathcal{D}_{a^+}^{\alpha;\varphi} [\varphi(t) - \varphi(a)]^{\beta-1} = \frac{\Gamma(\beta)}{\Gamma(\beta-\alpha)} [\varphi(t) - \varphi(a)]^{\beta-\alpha-1}$,
- ${}^c \mathcal{D}_{a^+}^{\alpha;\varphi} [\varphi(t) - \varphi(a)]^k = 0$, for all $k \in \{0, \dots, n-1\}$, $n \in \mathbb{N}$.

Lemma 4. Let $\alpha > 0$, $n \in \mathbb{N}$ such that $n-1 < q \leq n$. Then

- ${}^c \mathcal{D}_{a^+}^{q;\varphi} I_{a^+}^{\alpha;\varphi} u(t) = {}^c \mathcal{D}_{a^+}^{q-\alpha;\varphi} u(t)$; if $q > \alpha$.
- ${}^c \mathcal{D}_{a^+}^{q;\varphi} I_{a^+}^{\alpha;\varphi} u(t) u(t) = I_{a^+}^{\alpha-q;\varphi} u(t)$; if $\alpha > q$.

Lemma 5. Given a function $u \in C^n[a, b]$ and $0 < q < 1$, we have

$$|I_{a^+}^{q;\varphi} u(t_2) - I_{a^+}^{q;\varphi} u(t_1)| \leq \frac{2\|u\|}{\Gamma(q+1)} (\varphi(t_2) - \varphi(t_1))^q.$$

Finally, we recall the fixed point theorems that will be used to prove the main results. (We have C a Banach space in each theorem).

Lemma 6 (Banach fixed point theorem, [7]). Let U be a closed set in C and $\mathcal{T} : U \rightarrow U$ satisfies

$$|\mathcal{T}u - \mathcal{T}v| \leq \alpha|u - v|, \text{ for some } \alpha \in (0, 1), \text{ and for } u, v \in U.$$

Then \mathcal{T} admits one fixed point in U .

Lemma 7 (Krasnoselskii fixed point theorem, [15]). Let M be a closed, bounded, convex and nonempty subset of a Banach space U . Let A, B be operators such that

- (i) $Ax + By \in M$, where $x, y \in M$,
- (ii) A is compact and continuous,
- (iii) B is a contraction mapping.

Then there exists $z \in M$ such that $z = Az + Bz$.

Lemma 8 ([13]). For the p -Laplacian operator Ψ_p , the following conditions hold true:

- (1) If $|\delta_1|, |\delta_2| \geq \rho > 0$, $1 < p \leq 2$, $\delta_1 \delta_2 > 0$, then

$$|\Psi_p(\delta_1) - \Psi_p(\delta_2)| \leq (p-1)\rho^{p-2} |\delta_1 - \delta_2|.$$

- (2) If $p > 2$, $|\delta_1|, |\delta_2| \leq \rho_* > 0$, then

$$|\Psi_p(\delta_1) - \Psi_p(\delta_2)| \leq (p-1)\rho_*^{p-2} |\delta_1 - \delta_2|.$$

Lemma 9 ([10]). For nonnegative $a_i, i = 1, \dots, k$,

$$\left(\sum_{i=1}^k a_i \right)^q \leq k^{q-1} \left(\sum_{i=1}^k a_i^q \right), q \geq 1.$$

Now, we pass to prove the following result.

Lemma 10. For a given $h_m, g_m \in L^1(J, \mathbb{R}^3)$, the unique solution of the linear fractional initial value problem

$$\begin{cases} \mathcal{D}_{0^+}^{r_{1m}; \Phi} \Psi_p [\mathcal{D}_{0^+}^{r_{2m}; \Phi} (u_m(t) - I_{0^+}^{\sigma; \Phi} g_m(t))] = h_m(t), \\ m = 1, 2, 3, \text{ and } t \in J = (0, 1] \\ \Psi_p [\mathcal{D}_{0^+}^{r_{2m}; \Phi} (u_m(t) - I_{0^+}^{\sigma; \Phi} g_m(t))] |_{t=0} = 0, \\ u_m(0) = 0, \quad u_m(1) = \sum_{i=1}^n \lambda_{im} u_m(\zeta_{im}), \quad \zeta_{im} \in (0, 1] \\ \varphi(1) - \varphi(0) = K > 0. \end{cases}$$

is given by

$$\begin{aligned} u_m(t) &= \int_0^t \varphi_{r_{2m}}(t, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) h_m(e) de \right] ds + \int_0^t \varphi_{\sigma}(t, s) g_m(s) ds \quad (2.3) \\ &\quad - (\varphi(t) - \varphi(0)) \int_0^1 \varphi_{r_{2m}}(1, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) h_m(e) de \right] ds \\ &\quad + (\varphi(t) - \varphi(0)) \left(\sum_{i=1}^n \frac{\lambda_{im}}{K} u_m(\zeta_i) - \frac{g_m(0)}{K} \right). \end{aligned}$$

Proof. For $0 < r_{1m} < 1 < r_{2m} < 2$, Lemma 2 yields

$$\Psi_p [\mathcal{D}_{0^+}^{r_{2m}; \Phi} (u_m(t) - I_{0^+}^{\sigma; \Phi} g_m(t))] = I_{0^+}^{r_{1m}; \Phi} h_m(t) + c_{1m}$$

by conditions $\Psi_p [\mathcal{D}_{0^+}^{r_{2m}; \Phi} (u_m(t) - g_m(t))] |_{t=0} = 0$ we get $c_{1m} = 0$. Then

$$[\mathcal{D}_{0^+}^{r_{2m}; \Phi} (u_m(t) - I_{0^+}^{\sigma; \Phi} g_m(t))] = \Psi_q [I_{0^+}^{r_{1m}; \Phi} h_m(t)]$$

so

$$u_m(t) = I_{0^+}^{r_{2m}; \Phi} [\Psi_q [I_{0^+}^{r_{1m}; \Phi} h_m(t)]] + I_{0^+}^{\sigma; \Phi} g_m(t) + c_{2m} (\varphi(t) - \varphi(0)),$$

by conditions $u_m(0) = 0$ and $u_m(1) = \sum_{i=1}^n \lambda_{im} u_m(\zeta_i)$, we get

$$c_{2m} = \sum_{i=1}^n \frac{\lambda_{im}}{K} u_m(\zeta_i) - \frac{g_m(0)}{K} - I_{0^+}^{r_{2m}; \Phi} [\Psi_q [I_{0^+}^{r_{1m}; \Phi} h_m(t)]] |_{t=1}. \quad (2.4)$$

□

3. MAIN RESULTS

Taking into account Lemma 10, we define an operator $\mathcal{T} : \mathcal{C} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C} \times \mathcal{C}$

$$\mathcal{T}(u_1, u_2, u_3)(t) = (\mathcal{T}_1(u_1, u_2, u_3)(t), \mathcal{T}_2(u_1, u_2, u_3)(t), \mathcal{T}_3(u_1, u_2, u_3)(t)), \quad (3.1)$$

where (for $m = \overline{1, 3}$)

$$\begin{aligned} \mathcal{T}_m(u_1, u_2, u_3)(t) &= \int_0^t \varphi_{r_{2m}}(t, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \\ &\quad + \int_0^t \varphi_{\sigma}(t, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \end{aligned}$$

$$\begin{aligned}
 & -(\varphi(t) - \varphi(0)) \int_0^1 \varphi_{r_{2m}}(1, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \\
 & + (\varphi(t) - \varphi(0)) \left(\sum_{i=1}^n \frac{\lambda_{im}}{K} u_m(\zeta_i) - \frac{G_m(0, 0, 0, 0)}{K} \right), \tag{3.2}
 \end{aligned}$$

and

$$\begin{aligned}
 \varphi_{r_{2m}}(t, s) &= \frac{\varphi'(s) (\varphi(t) - \varphi(s))^{r_{2m}-1}}{\Gamma(r_{2m})}, & \varphi_{r_{1m}}(t, s) &= \frac{\varphi'(s) (\varphi(t) - \varphi(s))^{r_{1m}-1}}{\Gamma(r_{1m})}, \\
 \varphi_{\sigma}(t, s) &= \frac{\varphi'(s) (\varphi(t) - \varphi(s))^{\sigma-1}}{\Gamma(\sigma)}.
 \end{aligned}$$

For the sake of convenience, we use the following notations (for $m = \overline{1, 3}$):

$$\begin{aligned}
 \mathcal{K}_{1m} &= \frac{2^{q-2} (K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{k_m K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1}, \\
 \mathcal{K}_{2m} &= \left(\frac{K^{\sigma} l_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| \right), \\
 \mathcal{K}_{3m} &= \frac{2^{q-2} (K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{\mathcal{N} K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} + \frac{K^{\sigma} \mathcal{M}}{\Gamma(1+\sigma)} + \mathcal{M}, \\
 \mathcal{K}_{4m} &= \left(\frac{K^{r_{1m}} (\rho \mathcal{A}_m + \mathcal{N})}{\Gamma(1+r_{1m})} \right)^{q-2}, \\
 \mathcal{K}_{5m} &= \frac{(q-1)(1+K) K^{r_{2m}+r_{1m}} \mathcal{A}_m \mathcal{K}_{4m}}{\Gamma(1+r_{2m}) \Gamma(1+r_{1m})} + \frac{K^{\sigma} \mathcal{B}_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}|.
 \end{aligned}$$

3.1. An Existence and Uniqueness Result

Here, by using the Banach contraction mapping principle, we prove an existence and uniqueness result.

Theorem 1. Let $H_m, G_m : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ two continuous functions which satisfy the condition

(\mathbb{A}_1) there exist positive real constants $\mathcal{A}_m, \mathcal{B}_m$ such that, for all $t \in [0, 1]$ and $u_i, v_i \in \mathbb{R}, i, m = \overline{1, 3}$, we have

$$|H_m(t, u_1, u_2, u_3) - H_m(e, v_1, v_2, v_3)| \leq \mathcal{A}_m (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|),$$

$$|G_m(t, u_1, u_2, u_3) - G_m(e, v_1, v_2, v_3)| \leq \mathcal{B}_m (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|).$$

Then, system (1.1) admits a unique solution on $[0, 1]$ provided that

$$3(\mathcal{K}_{21} + \mathcal{K}_{22} + \mathcal{K}_{23}) < 1, \text{ and } \mathcal{K}_{51} + \mathcal{K}_{52} + \mathcal{K}_{53} < 1 \tag{3.3}$$

is valid.

Proof. We transform system (1.1) into a fixed point problem, $(u, v, w)(z) = \mathcal{T}(u, v, w)(z)$, where the operator \mathcal{T} is defined as in (3.1). Applying the Banach contraction mapping principle (Lemma 6), we show that the operator \mathcal{T} has a unique fixed point, which is the unique solution of system (1.1).

Let $\sup_{t \in [0,1]} H_m(t, 0, 0, 0) = \mathcal{N} < \infty$, and $\sup_{t \in [0,1]} G_m(t, 0, 0, 0) = \mathcal{M} < \infty$. Next, we set $\mathbb{U}\rho = \{(u_1, u_2, u_3) \in \mathcal{C} \times \mathcal{C} \times \mathcal{C}, \|(u_1, u_2, u_3)\| \leq \rho\}$, in which

$$\rho \geq \max \left\{ \sqrt[2-q]{3 \mathcal{K}_{1m}}, 3 \mathcal{K}_{3m}, m = \overline{1, 3} \right\}.$$

Observe that $\mathbb{U}\rho$ is a bounded, closed, and convex subset of \mathcal{C} . First, we show that $\mathcal{T}\mathbb{U}\rho \subset \mathbb{U}\rho$.

For any $(u, v, w) \in \mathbb{U}\rho$, $t \in [0, 1]$, using the condition (\mathbb{A}_1) , we have

$$\begin{aligned} |H_m(t, u, v, w)| &\leq |H_m(t, u, v, w) - H_m(e, 0, 0, 0)| + |H_m(e, 0, 0, 0)| \\ &\leq k_m(|u| + |v| + |w|) + N \leq \rho \mathcal{A}_m + \mathcal{N} \end{aligned}$$

and

$$|G_m(t, u, v, w)| \leq \rho \mathcal{B}_m + \mathcal{M}.$$

Then, we obtain

$$\begin{aligned} |\mathcal{T}_m(u_1, u_2, u_3)(t)| &\leq \left| \int_0^t \Phi_{r_{2m}}(t, s) \Psi_q \left[\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \right| \\ &\quad + \left| \int_0^t \Phi_{\sigma}(t, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \right| \\ &\quad + |(\varphi(t) - \varphi(0))| \left| \int_0^1 \Phi_{r_{2m}}(1, s) \Psi_q \left[\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \right| \\ &\quad + |(\varphi(t) - \varphi(0))| \left(\sum_{i=1}^n \frac{|\lambda_{im}|}{K} |u_m(\zeta_i)| + \frac{|G_m(0, 0, 0, 0)|}{K} \right), \end{aligned}$$

by Lemma 5 we get

$$\begin{aligned} |\mathcal{T}_m(u_1, u_2, u_3)(t)| &\leq \frac{(K+1)K^{r_{2m}}}{\Gamma(1+r_{2m})} \left| \Psi_q \left[\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] \right| \\ &\quad + \frac{K^{\sigma}(\rho \mathcal{B}_m + \mathcal{M})}{\Gamma(1+\sigma)} + \left(\sum_{i=1}^n \rho |\lambda_{im}| + \mathcal{M} \right), \end{aligned}$$

and by $\Psi_q(z) = |z|^{q-2}z$, we have

$$\begin{aligned} |\mathcal{T}_m(u_1, u_2, u_3)(t)| &\leq \frac{(K+1)K^{r_{2m}}}{\Gamma(1+r_{2m})} \left| \int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right|^{q-1} \\ &\quad + \frac{K^{\sigma}(\rho \mathcal{B}_m + \mathcal{M})}{\Gamma(1+\sigma)} + \left(\sum_{i=1}^n \rho |\lambda_{im}| + \mathcal{M} \right) \end{aligned}$$

$$\begin{aligned} &\leq \left(\frac{(\rho \mathcal{A}_m + \mathcal{N}) K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} \frac{(K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \\ &\quad + \frac{K^\sigma (\rho \mathcal{B}_m + \mathcal{M})}{\Gamma(1+\sigma)} + \left(\sum_{i=1}^n \rho |\lambda_{im}| + \mathcal{M} \right) \\ &\leq \frac{(K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} (\rho \mathcal{A}_m + \mathcal{N})^{q-1} \\ &\quad + \left(\frac{K^\sigma \mathcal{B}_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| \right) \rho + \frac{K^\sigma \mathcal{M}}{\Gamma(1+\sigma)} + \mathcal{M}. \end{aligned}$$

Thanks to Lemma 9, for all $m = \overline{1,3}$ we get

$$\begin{aligned} |\mathcal{T}_m(u_1, u_2, u_3)(t)| &\leq \frac{2^{q-2} (K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} (\mathcal{A}_m^{q-1} \rho^{q-1} + \mathcal{N}^{q-1}) \\ &\quad + \left(\frac{K^\sigma \mathcal{B}_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| \right) \rho + \frac{K^\sigma \mathcal{M}}{\Gamma(1+\sigma)} + \mathcal{M} \\ &\leq \frac{2^{q-2} (K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{\mathcal{A}_m K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} \rho^{q-1} \\ &\quad + \left(\frac{K^\sigma \mathcal{B}_m}{\Gamma(1+\sigma)} \sum_{i=1}^n |\lambda_{im}| \right) \rho + \frac{2^{q-2} (K+1) K^{r_{2m}}}{\Gamma(1+r_{2m})} \left(\frac{\mathcal{N} K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} \\ &\quad + \frac{K^\sigma \mathcal{M}}{\Gamma(1+\sigma)} + \mathcal{M} \\ &\leq \mathcal{K}_{1m} \rho^{q-1} + \mathcal{K}_{2m} \rho + \mathcal{K}_{3m} \leq \frac{\rho}{3}. \end{aligned}$$

Hence,

$$\|\mathcal{T}(u_1, u_2, u_3)\| \leq \sum_{m=1}^3 (\mathcal{K}_{1m} \rho^{q-1} + \mathcal{K}_{2m} \rho + \mathcal{K}_{3m}) \leq \rho, \tag{3.4}$$

which gives us $\mathcal{T}\mathbb{U}\sigma \subset \mathbb{U}\sigma$.

Next, we show that $\mathcal{T} : \mathcal{C} \times \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C} \times \mathcal{C}$ is a contraction.

Using condition (\mathbb{A}_1) , for any $(u_1, u_2, u_3), (v_1, v_2, v_3) \in \mathcal{C} \times \mathcal{C} \times \mathcal{C}$ and for each $t \in [0, 1]$, we have

$$\begin{aligned} &|\mathcal{T}_m(u_1, u_2, u_3) - \mathcal{T}_m(v_1, v_2, v_3)| \\ &\leq \left| \int_0^t \Phi_{r_{2m}}(t, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) ds \right. \\ &\quad \left. - \int_0^t \Phi_{r_{2m}}(t, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) ds \right| \end{aligned}$$

$$\begin{aligned}
& + \left| \int_0^t \Phi_{\sigma}(t, s) (G_m(s, u_1(s), u_2(s), u_3(s)) - G_m(s, v_1(s), v_2(s), v_3(s))) ds \right| \\
& + K \left| \int_0^1 \Phi_{r_{2m}}(1, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) ds \right. \\
& \left. - \int_0^1 \Phi_{r_{2m}}(1, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) ds \right| \\
& + \sum_{i=1}^n |\lambda_{im}| |u_m(\zeta_i) - v_m(\zeta_i)|,
\end{aligned}$$

by Lemma 5 and Lemma 8, we get

$$\begin{aligned}
& |\mathcal{T}_m(u_1, u_2, u_3) - \mathcal{T}_m(v_1, v_2, v_3)| \\
& \leq \frac{(1+K)K^{r_{2m}}}{\Gamma(1+r_{2m})} \left| \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) \right. \\
& \quad \left. - \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) \right| \\
& \quad + \left(\frac{K^{\sigma} \mathcal{B}_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| \right) (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|) \\
& \leq \frac{(1+K)K^{r_{2m}}(q-1)\mathcal{K}_{\mathcal{G}m}}{\Gamma(1+r_{2m})} \left| \int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right. \\
& \quad \left. - \int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right| \\
& \leq \left[\frac{(q-1)\mathcal{A}_m(1+K)K^{r_{2m}+r_{1m}}\mathcal{K}_{\mathcal{G}m}}{\Gamma(1+r_{2m})\Gamma(1+r_{1m})} + \frac{K^{\sigma}\mathcal{B}_m}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| \right] \\
& \quad \cdot (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|) \\
& \leq \mathcal{K}_{\mathcal{G}m} (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|).
\end{aligned}$$

Hence,

$$|\mathcal{T}(u_1, u_2, u_3) - \mathcal{T}_m(v_1, v_2, v_3)| \leq (\mathcal{K}_{\mathcal{S}1} + \mathcal{K}_{\mathcal{S}1} + \mathcal{K}_{\mathcal{S}3}) (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|).$$

Since $\mathcal{K}_{\mathcal{S}1} + \mathcal{K}_{\mathcal{S}1} + \mathcal{K}_{\mathcal{S}3} < 1$, by (3.3), the operator \mathcal{T} is a contraction. Therefore, using the Banach contraction mapping principle (Lemma 6), the operator \mathcal{T} has a unique fixed point. Hence, system (1.1) has a unique solution on $[0, 1]$. The proof is completed. \square

3.2. An Existence Result

Now we apply Krasnoselskii fixed point theorem (Lemma 7) to prove our second existence result. So, consider the following operator

$$\begin{aligned} \mathcal{T}(u_1, u_2, u_3)(t) &= \sum_{m=1}^3 (\mathcal{T}_m(u_1, u_2, u_3))(t) \\ &= \sum_{m=1}^3 (\mathcal{P}_{1m}(u_1, u_2, u_3)(t) + \sum_{m=1}^3 (\mathcal{P}_{2m}(u_1, u_2, u_3)(t) \\ &= \mathcal{P}_1(u_1, u_2, u_3)(t) + \mathcal{P}_2(u_1, u_2, u_3)(t), \end{aligned}$$

where

$$\begin{aligned} \mathcal{P}_1(u_1, u_2, u_3)(t) &= \sum_{m=1}^3 (\mathcal{P}_{1m}(u_1, u_2, u_3)(t), \\ \mathcal{P}_2(u_1, u_2, u_3)(t) &= \sum_{m=1}^3 (\mathcal{P}_{2m}(u_1, u_2, u_3)(t), \end{aligned}$$

and

$$\begin{aligned} (\mathcal{P}_{1m}(u_1, u_2, u_3)(t) &= \int_0^t \varphi_{r_{2m}}(t, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \\ &- (\varphi(t) - \varphi(0)) \int_0^1 \varphi_{r_{2m}}(1, s) \Psi_q \left[\int_0^s \varphi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right] ds \end{aligned}$$

and

$$\begin{aligned} (\mathcal{P}_{2m}(u_1, u_2, u_3)(t) &= \int_0^t \varphi_{\sigma}(t, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \\ &+ (\varphi(t) - \varphi(0)) \left(\sum_{i=1}^n \frac{\lambda_{im}}{K} u_m(\zeta_i) - \frac{G_m(0, u_1(0), u_2(0), u_3(0))}{K} \right). \end{aligned}$$

Theorem 2. Let $H_m, G_m : [0, 1] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ be continuous functions which satisfy condition (A_1) in Theorem 1.

In addition, we assume that there exist two positive constants $\Upsilon_{1m}, \Upsilon_{2m}$ such that, for all $t \in [0, 1]$ and $u_i, v_i \in \mathbb{R}, i, m = \overline{1, 3}$, we have

$$\begin{aligned} |H_m(t, u_1, u_2, u_3)| &\leq \Upsilon_{1m}, \\ |G_m(t, u_1, u_2, u_3)| &\leq \Upsilon_{2m}. \end{aligned}$$

Moreover, assume that

$$\sum_{i=1}^n |\lambda_{im}| \leq \frac{1}{3}, \text{ and } \left(\sum_{m=1}^3 \frac{(q-1) \mathcal{A}_m (1+K) K^{r_{2m}+r_{1m}} \mathcal{K}_{\mathcal{A}_m}}{\Gamma(1+r_{2m}) \Gamma(1+r_{1m})} \right) < 1.$$

Then, problem (1.1) admits at least one solution on $[0, 1]$.

Proof. The proof will be given in several steps. Let

$$\mathbb{U}_\delta = \{(u_1, u_2, u_3) \in \mathcal{C} \times \mathcal{C} \times \mathcal{C}, \|(u_1, u_2, u_3)\| \leq \delta\},$$

in which

$$\delta \geq 3 \max_{m=1,3} \left[\frac{\left(\frac{\Upsilon_{1m} K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} \frac{(K+1)K^{r_{2m}}}{\Gamma(1+r_{2m})} + \frac{K^\sigma \Upsilon_{2m}}{\Gamma(1+\sigma)} + \mathcal{M}}{\frac{1}{3} - \sum_{i=1}^n |\lambda_{im}|} \right].$$

Step 1. We prove that

$$|(\mathcal{T}(u_1, u_2, u_3))(t)| \leq \delta.$$

Let $(u_1, u_2, u_3), (v_1, v_2, v_3), (w_1, w_2, w_3) \in \mathbb{U}_\delta$. As in the proof of Theorem 1, we have

$$\begin{aligned} & |(\mathcal{P}_{1m}(u_1, u_2, u_3)(t) + \mathcal{P}_{2m}(u_1, u_2, u_3)(t))| \\ & \leq \left(\frac{\Upsilon_{1m} K^{r_{1m}}}{\Gamma(1+r_{1m})} \right)^{q-1} \frac{(K+1)K^{r_{2m}}}{\Gamma(1+r_{2m})} + \frac{K^\sigma \Upsilon_{2m}}{\Gamma(1+\sigma)} + \mathcal{M} + \left(\sum_{i=1}^n |\lambda_{im}| \right) \delta \leq \frac{\delta}{3}. \end{aligned}$$

Hence

$$\begin{aligned} |(\mathcal{T}(u_1, u_2, u_3))(t)| & = |(\mathcal{T}_1(u_1, u_2, u_3))(t) + (\mathcal{T}_2(u_1, u_2, u_3))(t) + (\mathcal{T}_3(u_1, u_2, u_3))(t)| \\ & \leq |(\mathcal{T}_1(u_1, u_2, u_3))(t)| + |(\mathcal{T}_2(u_1, u_2, u_3))(t)| + |(\mathcal{T}_3(u_1, u_2, u_3))(t)| \\ & \leq \sum_{m=1}^3 |\mathcal{P}_{1m}(u_1, u_2, u_3)(t) + \mathcal{P}_{2m}(u_1, u_2, u_3)(t)| \leq \delta. \end{aligned}$$

Accordingly, $\sum_{m=1}^3 \mathcal{P}_{1m}(u_1, u_2, u_3)(t) + \mathcal{P}_{2m}(u_1, u_2, u_3)(t) \in \mathbb{U}_\delta$ and the condition (i) of Lemma 7 is satisfied.

Step 2. \mathcal{P}_1 is a contraction.

Let $(u_1, u_2, u_3), (v_1, v_2, v_3) \in \mathbb{U}_\delta$. We have the following estimate

$$\begin{aligned} & |\mathcal{P}_{1m}(u_1, u_2, u_3)(t) - \mathcal{P}_{1m}(v_1, v_2, v_3)(t)| \\ & \leq \left| \int_0^t \Phi_{r_{2m}}(t, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) ds \right. \\ & \quad \left. - \int_0^t \Phi_{r_{2m}}(t, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) ds \right| \\ & \quad + K \left| \int_0^1 \Phi_{r_{2m}}(1, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) ds \right. \\ & \quad \left. - \int_0^1 \Phi_{r_{2m}}(1, s) \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) ds \right| \\ & \leq \frac{(1+K)K^{r_{2m}}}{\Gamma(1+r_{2m})} \left| \Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, u_1(e), u_2(e), u_3(e)) de \right) \right| \end{aligned}$$

$$\begin{aligned} & -\Psi_q \left(\int_0^s \Phi_{r_{1m}}(s, e) H_m(e, v_1(e), v_2(e), v_3(e)) de \right) \\ & \leq \frac{(q-1) \mathcal{A}_m (1+K) K^{r_{2m}+r_{1m}} \mathcal{K}_{4m}}{\Gamma(1+r_{2m}) \Gamma(1+r_{1m})} (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|). \end{aligned}$$

So

$$\begin{aligned} & | \mathcal{P}_1(u_1, u_2, u_3)(t) - \mathcal{P}_1(v_1, v_2, v_3)(t) | \\ & \leq \left(\sum_{m=1}^3 \frac{(q-1) \mathcal{A}_m (1+K) K^{r_{2m}+r_{1m}} \mathcal{K}_{4m}}{\Gamma(1+r_{2m}) \Gamma(1+r_{1m})} \right) (|u_1 - v_1| + |u_2 - v_2| + |u_3 - v_3|). \end{aligned}$$

Since $\left(\sum_{m=1}^3 \frac{(q-1) \mathcal{A}_m (1+K) K^{r_{2m}+r_{1m}} \mathcal{K}_{4m}}{\Gamma(1+r_{2m}) \Gamma(1+r_{1m})} \right) < 1$, the operator \mathcal{P}_1 is a contraction.

Step 3. \mathcal{P}_2 is compact and continuous.

Since H_m, G_m are a continuous functions, this implies that the operator \mathcal{P}_2 is continuous on \mathbb{U}_δ . Moreover, $\mathcal{P}_2(u_1, u, u_3)$ is uniformly bounded by (3.4). Next, we show equicontinuity. Let $(u_1, u, u_3) \in \mathbb{U}_\delta$, we have

$$\begin{aligned} |(\mathcal{P}_{2m}(u_1, u_2, u_3)(t))| & \leq \left| \int_0^t \Phi_\sigma(t, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \right| \\ & \quad + K \left(\sum_{i=1}^n \frac{|\lambda_{im}|}{K} |u_m(\zeta_i)| + \frac{|G_m(0, u_1(0), u_2(0), u_3(0))|}{K} \right) \\ & \leq \left(\frac{\Upsilon_{2m} K^\sigma}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| + \Upsilon_{2m} \right) \delta. \end{aligned}$$

So

$$|(\mathcal{P}_2(u_1, u_2, u_3)(t))| \leq \sum_{m=1}^3 \left(\frac{\Upsilon_{2m} K^\sigma}{\Gamma(1+\sigma)} + \sum_{i=1}^n |\lambda_{im}| + \Upsilon_{2m} \right) \delta. \tag{3.5}$$

Moreover, $\mathcal{P}_2(u_1, u, u_3)$ is uniformly bounded by (3.5). Next, we show equicontinuity and $t_1, t_2 \in [0, 1]$ such that $t_1 < t_2$ we have

$$\begin{aligned} & |(\mathcal{P}_{2m}(u_1, u_2, u_3)(t_2) - (\mathcal{P}_{2m}(u_1, u_2, u_3)(t_1))| \\ & \leq \left| \int_0^{t_2} \Phi_\sigma(t_2, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds - \int_0^{t_1} \Phi_\sigma(t_1, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \right| \\ & \leq \frac{\Upsilon_{2m}}{\Gamma(1+\sigma)} (\varphi(t_2) - \varphi(t_1))^\sigma. \end{aligned}$$

So

$$\begin{aligned} & |(\mathcal{P}_2(u_1, u_2, u_3)(t_2) - (\mathcal{P}_2(u_1, u_2, u_3)(t_1))| \\ & \leq \left| \int_0^{t_2} \Phi_\sigma(t_2, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds - \int_0^{t_1} \Phi_\sigma(t_1, s) G_m(s, u_1(s), u_2(s), u_3(s)) ds \right| \end{aligned}$$

$$\leq \sum_{m=1}^3 \left(\frac{\Upsilon_{2m}}{\Gamma(1+\sigma)} \right) (\varphi(t_2) - \varphi(t_1))^\sigma.$$

Consequently,

$$|(\mathcal{P}_2(u_1, u_2, u_3)(t_2) - \mathcal{P}_2(u_1, u_2, u_3)(t_1))| \rightarrow 0, \text{ as } t_1 \rightarrow t_2.$$

This shows that $\mathcal{P}_2\mathbb{U}_\delta$ is equicontinuous. Hence, by Arzelià-Ascoli theorem \mathcal{P}_2 is completely continuous on \mathbb{U}_δ . As a consequence of Krasnoselskii's fixed point theorem, we conclude that it has a fixed point which is a solution of (1.1). The proof of Theorem 2 is thus completely achieved. \square

4. AN ILLUSTRATIVE EXAMPLE

Consider the following nonlinear equation for all $t \in (0, 1]$:

$$\left\{ \begin{array}{l} {}^c \mathcal{D}_{0^+}^{\frac{1}{2}; t^2} \Psi_p \left[{}^c \mathcal{D}_{0^+}^{\frac{1}{4}; t^2} \left(u(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{e^t + 1}{1 + (u(t))^2} \right) \right) \right] = \frac{1}{1+t^2} \left(\frac{u(t)}{1 + (u(t))^2} \right), \\ {}^c \mathcal{D}_{0^+}^{\frac{1}{4}; t^2} \Psi_p \left[{}^c \mathcal{D}_{0^+}^{\frac{1}{2}; t^2} \left(v(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{t^2 + 1}{1 + (v(t))^2} \right) \right) \right] = \frac{t}{1+e^t} \left(\frac{v(t)}{1 + (v(t))^2} \right), \\ {}^c \mathcal{D}_{0^+}^{\frac{3}{4}; t^2} \Psi_p \left[{}^c \mathcal{D}_{0^+}^{\frac{3}{4}; t^2} \left(w(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{e^t}{1 + (w(t))^2} \right) \right) \right] = \frac{e^t}{1+t^2} \left(\frac{w(t)}{1 + (w(t))^2} \right), \\ \Psi_p \left[{}^c \mathcal{D}_{0^+}^{r_2; \varphi} \left(u(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{e^t + 1}{1 + (u(t))^2} \right) \right) \right] \Big|_{t=0} = 0, \\ \Psi_p \left[{}^c \mathcal{D}_{0^+}^{r_2; \varphi} \left(v(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{t^2 + 1}{1 + (v(t))^2} \right) \right) \right] \Big|_{t=0} = 0, \\ \Psi_p \left[{}^c \mathcal{D}_{0^+}^{r_2; \varphi} \left(w(t) - I_{0^+}^{\sigma; \varphi} \left(\frac{e^t}{1 + (w(t))^2} \right) \right) \right] \Big|_{t=0} = 0, \\ u(0) = v(0) = w(0) = 0, \\ u(1) = \sum_{i=1}^n \frac{1}{7(i!)} u(\zeta_i), v(1) = \sum_{i=1}^n \frac{1}{9(i!)} v(\zeta_i), w(1) = \sum_{i=1}^n \frac{1}{11(i!)} w(\zeta_i), \zeta_i \in (0, 1] \end{array} \right. \quad (4.1)$$

and

$$\begin{aligned} K &= 1, \\ \Upsilon_{11} &= \mathcal{A}_1 = \frac{1}{2}, \Upsilon_{12} = \mathcal{A}_2 = \frac{1}{4}, \Upsilon_{13} = \mathcal{A}_3 = \frac{e}{2}, \\ \Upsilon_{21} &= \mathcal{B}_1 = \frac{1+e}{2}, \Upsilon_{22} = \mathcal{B}_2 = 1, \Upsilon_{23} = \mathcal{B}_3 = \frac{e}{2}. \end{aligned}$$

Thus, the assumptions (\mathbb{A}_1) are satisfied and Theorem 1-2 implies that (4.1) has a unique solution on $[0, 1]$.

REFERENCES

- [1] H. Beddani and Z. Dahmani, "Solvability for nonlinear differential problem of Langevin type via phi-Caputo approach." *Eur. J. Math.Appl.*, vol. 1, no. 2, pp. 1–11, 2021, doi: [10.28919/ejma.2021.1.11](https://doi.org/10.28919/ejma.2021.1.11).
- [2] A. Devi, A. Kumar, D. Baleanu, and A. Khan, "On stability analysis and existence of positive solutions for a general non-linear fractional differential equations." *Advances in Difference Equations.*, vol. 300, 2020, doi: [10.1186/s13662-020-02729-3](https://doi.org/10.1186/s13662-020-02729-3).
- [3] A. Aghajani, E. Pourhadi, and J. J. Trujillo, "Application of measure of noncompactness to a Cauchy problem for fractional differential equations in Banach spaces." *Fract. Calc. Appl. Anal.*, vol. 16, pp. 962–977, 2013, doi: [10.2478/s13540-013-0059-y](https://doi.org/10.2478/s13540-013-0059-y).
- [4] O. P. Agrawal, "Some generalized fractional calculus operators and their applications in integral equations." *Fract. Calc. Appl. Anal.*, vol. 15, pp. 700–711, 2012, doi: [10.2478/s13540-012-0047-7](https://doi.org/10.2478/s13540-012-0047-7).
- [5] B. Ahmad, M. M. Matar, and O. M. EL-Salmy, "Existence of solutions and Ulam stability for Caputo type sequential fractional differential." *Int. J. Anal. Appl.*, vol. 15, pp. 86–101, 2017.
- [6] R. Almeida, "A Caputo fractional derivative of a function with respect to another function." *Mathematics. Commun. Nonlinear Sci. Numer. Simul.*, vol. 44, pp. 460–481, 2017, doi: [10.1016/j.cnsns.2016.09.006](https://doi.org/10.1016/j.cnsns.2016.09.006).
- [7] K. Deimling, *Nonlinear Functional Analysis*. SpringerLink., 1985. doi: [10.1007/978-3-662-00547-7](https://doi.org/10.1007/978-3-662-00547-7).
- [8] K. Diethelm, *The Analysis of Fractional Differential Equations*. SpringerLink., 2010.
- [9] S. Etemad, M. M. Matar, M. A. Ragusa, and S. Rezapour, "Tripled fixed points and existence study to a tripled impulsive fractional differential system via measures of noncompactness." *Mathematics.*, vol. 25, no. 10, pp. 1–17, 2022, doi: [10.3390/math10010025](https://doi.org/10.3390/math10010025).
- [10] R. Herrmann, *Fractional Calculus. An Introduction for Physicists*. World Scientific Publ., March 2014. doi: [10.1142/8934](https://doi.org/10.1142/8934).
- [11] M. D. Kassim and N. E. Tatar, "Stability of logarithmic type for a Hadamard fractional differential problem." *J. Pseudo-Differ. Oper. Appl.*, vol. 11, pp. 447–466, 2020.
- [12] H. Khan, T. Abdeljawad, M. Aslam, R. A. Khan, and A. Khan, "Existence of positive solution and Hyers-Ulam stability for a nonlinear singular-delay-fractional differential equation." *Advances in Difference Equations.*, vol. 104, 2019, doi: [10.1186/s13662-019-2054-z](https://doi.org/10.1186/s13662-019-2054-z).
- [13] H. Khan, W. Chen, and H. Sun, "Analysis of positive solution and Hyers-Ulam stability for a class of singular fractional differential equations with p-Laplacian in Banach space." *Math. Methods Appl. Sci.*, vol. 41, no. 9, pp. 3430–3440, 2018, doi: [10.1002/mma.4835](https://doi.org/10.1002/mma.4835).
- [14] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*. Elsevier Science., 2006.
- [15] M. A. Krasnoselskii's, "Two remarks on the method of successive approximations." , vol. 10, no. 1, pp. 123–127, 1955.
- [16] Y. Li, "Existence of positive solutions for fractional differential equation involving integral boundary conditions with p -Laplacian operator." *Adv. Differ. Equ.*, vol. 1, no. 135, 2017.
- [17] A. Mahdjouba, J. J. Nieto, and A. Ouahab, "System of fractional boundary value problem with p-Laplacian and advanced arguments." *Advances in Difference Equations.*, vol. 352, 2021, doi: [10.1186/s13662-021-03508-4](https://doi.org/10.1186/s13662-021-03508-4).
- [18] T. J. Osler, "Fractional derivatives of a composite function." *SIAMJ Math Anal.*, vol. 1, no. 2, pp. 288–293, 1970, doi: [10.1137/0501026](https://doi.org/10.1137/0501026).
- [19] I. Podlubny, *Fractional Differential Equations*, Academic Press. Academic Press, 1999.
- [20] S. G. Samko, A. A. Kilbas, and O. I. Mariche, "Fractional integrals and derivatives." *translated from the 1987 Russian original. Yverdon.*, 1993.

*Authors' addresses***Hamid Beddani**

(**Corresponding author**) Higher School of Electrical and Energy Engineering, Laboratory of Complex Systems, Oran, Algeria

E-mail address: beddanihamid@gmail.com

Moustafa Beddani

E. N. S. of Mostaganem, Department of Exact Sciences, Mostaganem, Algeria

E-mail address: beddani2004@yahoo.fr

Zoubir Dahmani

Laboratory of Pure and Applied Mathematics, Abdelhamid Bni Badis University, 27000, Algeria

E-mail address: zzdahmani@yahoo.fr